

What is the fate of the heavy electron at a quantum critical point?

¹ P. Coleman and ² C. Pépin

¹Center for Materials Theory, Department of Physics and Astronomy, Rutgers University, Piscataway, NJ 08854, USA.

²SPhT, L'Orme des Merisiers, CEA-Saclay, 91191 Gif-sur-Yvette, France.

A growing body of evidence suggests that the quantum critical behavior at the onset of magnetism in heavy fermion systems can not be understood in terms of a simple quantum spin density wave. This talk will discuss the consequences of this conclusion, touching on its possible implications in the realm of two dimensional systems and outlining current theoretical and experimental efforts to characterize the nature of the critical point in heavy fermion materials.

PACS No:

A. Introduction.

Discoveries over the past decade have brought a new awareness of the importance of quantum critical points in condensed matter physics. A quantum critical point (QCP) is a zero-temperature instability between two phases of matter where quantum fluctuations develop long range correlations in both space and time [1]. These special points exert wide-reaching influence on the finite temperature properties of a material. Systems close to quantum criticality develop a new excitation structure, they display novel thermodynamic, transport and magnetic behavior. They also have marked a predeliction towards the development of new kinds of order, such as anisotropic superconductivity. A dramatic example is provided by the cuprate superconductors. By doping with holes, these materials pass through one or more quantum phase transitions: from a Mott insulator to a




FIG. 1. Schematic phase diagram for cuprate superconductors showing location of possible quantum critical points. One of these QCP may be responsible for the anomalous normal state which develops above the pseudogap scale.

metal with a spin gap at low doping, and at higher doping a second QPT may occur when the spin gap closes [2] (Fig. 1). The singular interactions induced by quantum criticality may be the driving force for both the high temperature superconductivity and the anomalous metallic state above the spin gap temperature T^* . [3]

This paper discusses quantum criticality in heavy fermion materials. These materials offer a unique opportunity to study quantum criticality under controlled conditions. By the application of pressure or doping, heavy fermion materials can be tuned through a quantum critical point from a metallic antiferromagnet into a paramagnet (Fig. 2). Unlike the cuprate metals, here the paramagnetic phase is a well characterized Fermi liquid, [4–6] with heavy Landau quasiparticles, or “heavy electrons”. A central property of these quasiparticles, is the existence of a finite overlap “ Z ” between a single quasiparticle state, denoted by $|qp^- \rangle$ and the state formed by adding a single electron to the ground-state, denoted by $|e^- \rangle = c_{\mathbf{k}\sigma}^\dagger |0 \rangle$. This quantity is closely related to the ratio m/m^* of the electron to quasiparticle mass,

$$Z = |\langle e^- | qp^- \rangle|^2 \sim \frac{m}{m^*}. \quad (1)$$

A wide body of evidence suggests that m^*/m diverges at a heavy fermion QCP, indicating that

$$Z \rightarrow 0 \quad (P \rightarrow P_c).$$

The state which forms at the QCP is referred to as a “non-Fermi” or “singular Fermi liquid”. [7,8] By what mechanism does this break-down in the Landau quasiparticle occur?

B. Properties of the Heavy Fermion Quantum Critical Point

There is a growing list of heavy fermion systems that have been tuned to an antiferromagnetic QCP by the application of pressure or by doping (Table 1.). These materials display many common properties

Table. 1. Selected Heavy Fermion compounds with quantum critical points.

Compound	$x_c/P_c/H_c$	$\frac{C_v}{T} \rightarrow \infty?$	$\rho \sim T^a$	Ref.
$YbRh_2Si_{2-x}Ge_x$	$x_c = 0.1$	$\text{Log} \left(\frac{T_0}{T} \right)^{(b)}$	T	[9]
$CeCu_{6-x}Au_x$	$x_c = 0.1$	$\text{Log} \left(\frac{T_0}{T} \right)$	$T + c$	[10–12]
$CeCu_{6-x}Ag_x$	$x_c = 0.09$	$\text{Log} \left(\frac{T_0}{T} \right)$	$T^{1.1}$	[13]
$CeNi_2Ge_2$	$P_c = 0$	$\text{Log} \left(\frac{T_0}{T} \right)^{(a)}$	$T^{1.2}$	[14–18]
U_2Pt_2In	$P_c = 0$	$\text{Log} \left(\frac{T_0}{T} \right)$	T	[19]
U_2Pd_2In	$P_c < 0$	?	$T + c$	[19]
$CePd_2Si_2$	$P_c > 0$	?	$T^{1.2}$	[14]
$CeIn_3$	$P_c > 0$	?	$T^{1.5}$	[14,20]
$U_3Ni_3Sn_4$	$P_c > 0$	no	?	[21]
$CeCu_2Si_2$	$P_c = 0$	no	$T^{1.5}$	[22]
$CeRu_2Si_2$	$H_c^ = 7.7T$	$\text{Log} \left(\frac{T_0}{T} \right)$		[23]
UPt_3	$H_c^ = 20T$	$\text{Log} \left(\frac{T_0}{T} \right)$	$T^{1.2}$	[24]
$Sr_3Ru_2O_7$	$H_c^ = 7.7T$	$\text{Log} \left(\frac{T_0}{T} \right)$	$T^{1.1}$	[25]

^(a) New data [18] show a stronger divergence at lower temperatures, and $\gamma \sim A - B\sqrt{T}$ at intermediate temperatures.

^(b) At low temperatures, γ diverges more rapidly than $\text{Log} \left(\frac{T}{T_0} \right)$ [9].

- **Fermi liquid behavior in the paramagnet**, as indicated by the emergence of a quadratic temperature dependence in the resistivity in the approach to the QPT $\rho = \rho_0 + AT^2$ [21,26] at ever lower temperatures.
- **Divergent specific heat** at the QCP, typically with a logarithmic temperature dependence,

$$\gamma(T) = \frac{C_v(T)}{T} = \gamma_0 \log \left[\frac{T_0}{T} \right], \quad (2)$$

suggesting that the Fermi temperature vanishes and the quasiparticle effective masses diverge

$$T_F^* \rightarrow 0, \quad \frac{m^*}{m} \rightarrow \infty \quad (3)$$

at the QCP. Further support for this conclusion is provided by the observation that the quadratic coefficient A of the resistivity grows, and may diverge in the approach to the quantum critical point [19].

- **Quasi-linear resistivity**

$$\rho \propto T^{1+\epsilon}, \quad (4)$$

at the QCP with ϵ in the range $0 - 0.6$. In critical $YbRh_2Si_{2-x}Ge_x$, $\rho \propto T$ over three decades [9].

- **Non-Curie spin susceptibilities**

$$\chi^{-1}(T) = \chi_0^{-1} + cT^a \quad (5)$$

with $a < 1$ observed in critical $CeCu_{6-x}Au_x$ ($x=0.1$), $YbRh_2Si_{2-x}Ge_x$ ($x=0.1$) and $CeNi_2Ge_2$.

- **E/T and H/T Scaling.** In critical $CeCu_{6-x}Au_x$ and $YbRh_2Si_{2-x}Ge_x$ the differential magnetic susceptibility dM/dH exhibits H/T scaling,

$$(dM/dH)^{-1} = \chi_0^{-1} + cT^a g[H/T], \quad (6)$$

where $a \approx 0.75$. Neutron measurements [12] show E/T scaling [27,28] in the dynamical spin susceptibility of critical $CeCu_{6-x}Au_x$, throughout the Brillouin zone, parameterized in the form

$$\chi^{-1}(\mathbf{q}, \omega) = T^a f(E/T) + \chi_0^{-1}(\mathbf{q}) \quad (7)$$

$F[x] \propto (1 - ix)^a$. Scaling behavior with a single anomalous exponent in the momentum-independent component of the dynamical spin susceptibility suggests an emergence of *local* magnetic moments which are *critically correlated in time* at the quantum critical point [12].




FIG. 2. Illustration of quantum critical physics in heavy fermion metals. As criticality is approached from either side of the transition, the temperature scale T_F^* on which Fermi liquid behavior breaks down goes to zero. A key challenge is to characterize the new class of universal excitations which develops above T_F^* .

Growing evidence suggests a strong connection between the heavy fermion QCP and “metamagnetism”. A number of strongly correlated electron systems, such as $CeRu_2Si_2$ [23] and UPt_3 [24], and $Sr_3Ru_2O_7$ [25] exhibit a sudden rise in the magnetic polarization at a finite “critical” field. Properties characteristic of an heavy fermion QCP, (such as the logarithmic dependence of the specific heat and a quasi-linear resistivity) are seen to develop at such metamagnetic transitions. In all likelihood, this reflects the close vicinity to a *ferromagnetic* QCP (Fig 3), but why is the physics similar to that of a heavy fermion QCP?




FIG. 3. Schematic phase diagram for a metamagnet. Second order lines radiate from the tricritical point[29], intersecting the $T = 0$ plane at quantum critical points. Ambient pressure is denoted by a dotted line. When the material passes close to the QCP, a metamagnetic transition takes place.

C. Universality

Usually, the physics of a metal above its Fermi temperature depends on the detailed chemistry and band-structure of the material: it is non-universal. However, if the renormalized Fermi temperature $T_F^*(P)$ can be tuned to become arbitrarily small compared with the characteristic scales of the material as one approaches a QCP, we expect that the “high energy” physics *above* the Fermi temperature T_F^* is itself, universal.

Quantum critical behavior implies a divergence of the long distance and long-time correlations in the material. Finite temperatures introduce the cutoff timescale

$$\tau_T = \frac{\hbar}{k_B T} \quad (8)$$

beyond which coherent quantum processes are dephased by thermal fluctuations. Renormalization group principles [30] imply that the quantum critical physics has an upper-critical dimension d_u . For $d < d_u$, the τ_T becomes the correlation time τ of the system [31], so frequency dependent correlation functions and response functions take the form

$$F(\omega, T) = \frac{1}{\omega^\alpha} f(\omega \tau_T) = \frac{1}{\omega^\alpha} f(\hbar \omega / k_B T). \quad (9)$$

leading to E/T scaling [32]. By contrast, for $d > d_u$ the correlation time is sensitive to the details of the short-distance interactions between the critical modes, and in general $\tau^{-1} \propto T^{1+b}$. Thus E/T scaling with a non-trivial

exponent strongly suggests that the underlying physics of the heavy fermion quantum critical point is governed by universal physics with $d_u > 3$.

D. Failure of the Spin Density Wave picture

The commonly accepted picture of the heavy fermion QCP assumes the non-Fermi liquid behavior derives from Bragg diffraction of the electrons off a quantum-critical spin density wave (QSDW) [30,33–35]. The virtual emission of these soft fluctuations,

$$e^- \rightleftharpoons e^- + \text{spin fluctuation} \quad (10)$$

generates a retarded interaction

$$V_{eff}(\mathbf{q}, \omega) = g^2 \left[\frac{\chi(\mathbf{q}, \omega)}{(\mathbf{q} - \mathbf{Q})^2 + \xi^{-2} - \frac{i\omega}{\Gamma_{\mathbf{Q}}}} \right] \quad (11)$$

between the electrons, where $\chi(\mathbf{q}, \omega)$ is the dynamical spin susceptibility of the collective modes. The damping term $-i\omega/\Gamma_{\mathbf{Q}}$ of the magnetic fluctuations is derived from the linear density of particle-hole states in the Fermi sea. ξ^{-1} and $\tau^{-1} = \Gamma_{\mathbf{Q}} \xi^{-2}$ are the inverse spin correlation length and correlation times respectively. In real space,

$$V_{eff}(r, \omega = 0) \propto \frac{e^{-r/\xi}}{r} e^{i\mathbf{Q} \cdot \mathbf{r}} \quad (12)$$

is a “modulated” Yukawa potential whose range $\xi \sim (P - P_c)^{-\frac{1}{2}} \rightarrow \infty$ at the QCP. Unlike a ferromagnetic QCP, the modulated potential only effects electron quasiparticles along “hot lines” on the Fermi surface, that are separated by the wave-vector $\mathbf{Q}$ and satisfy $\epsilon_{\mathbf{k}} = \epsilon_{\mathbf{k}+\mathbf{Q}}$. At a finite temperature, electrons within a momentum range $\sim \sqrt{T}$ are affected by this critical scattering (Fig. 4.). This limits the ability of this singular potential to generate non Fermi liquid behavior.




FIG. 4. Quantum spin density wave scenario, where the Fermi surface “folds” along lines separated by the magnetic $\mathbf{Q}$ vector, pinching off into two separate Fermi surface sheets.

There are then two major difficulties with the QSDW scenario for the heavy fermion QCP:

1. **No breakdown of the Fermi liquid** Away from the hot lines, the Fermi surface and Landau quasiparticles remain intact at the QCP. Thus the specific heat and typical quasiparticle mass do not diverge but exhibit a weaker singularity, $C_V/T = \gamma_o - A\sqrt{T}$ in the QSDW picture [35].
2. **No E/T scaling** The quantum critical behavior predicted by this model has been extensively studied [30,35]. In the interaction $V_{eff}(\mathbf{q}, \omega)$ the momentum dependence enters with twice the power of the frequency, so

$$\tau \sim \xi^z, \quad (z = 2).$$

In the renormalization group (RG) treatment [30] time counts as z space dimensions so the effective dimensionality is $D_{eff} = d + z = d + 2$. The upper critical dimension is set by $D_{eff} = 4$, or $d_u = 2$ [35], so 3D quantum spin fluctuations will not lead to E/T scaling. In three dimensions, QSDW theory predicts that the scale entering into the energy dependent response functions should scale as $T^{3/2}$, with a non-universal prefactor [1].

1. Can a 2D spin fluid cure the difficulties?

One explanation of E/T scaling and the logarithmically divergent specific heat [36] is to suppose that the spin fluctuations form a quasi-two-dimensional spin fluid [36,20], lying at the critical dimension. Inelastic neutron scattering experiments on $CeCu_{6-x}Au_x$, ($x=0.1$) support a kind of reduced dimensionality in which the critical scattering is concentrated along linear, rather than at point-like regions in reciprocal space [12,36]. More recent data [37] may support quasi-2D spin fluctuations at intermediate scales in $CeGe_2Ni_2$.

E. New Approaches

Unfortunately, quasi-two dimensionality still cannot explain the anomalous exponents in the E/T scaling, and we are essentially forced to consider the possibility of a fundamentally new *interacting* fixed point. Two approaches have emerged:

1. Local Quantum Criticality

The momentum-independent scaling term in the inverse dynamic susceptibility (7) suggests that the critical

behavior associated with the heavy fermion QCP contains some kind of *local* critical excitation [12]. One possibility, is that this critical excitation is the spin itself, which would then presumably develop a slow power-law decay [32,38]

$$\langle S(\tau)S(\tau') \rangle = \frac{1}{(\tau - \tau')^{2-\epsilon}}, \quad (13)$$

where $\epsilon \neq 0$ signals non-Fermi liquid behavior.




FIG. 5. In the local quantum critical theory, each spin behaves as a local moment in a fluctuating Weiss field. In the theory of Si et al [39], a self-consistent solution can be obtained for $\epsilon = 1$ in which the local susceptibility develops a self-energy with a non-universal exponent. $M(\omega) \propto \omega^{\frac{1}{\rho\Lambda}}$.

Si [39] et al have developed this idea, proposing that the *local* spin susceptibility $\chi_{loc} = \sum_{\vec{q}} \chi(\vec{q}, \omega)|_{\omega=0}$ diverges at a heavy fermion QCP, From (7),

$$\chi_{loc}(T) \sim \int d^d q \frac{1}{(\mathbf{q} - \mathbf{Q})^2 + T^\alpha} \sim T^{(d-2)\alpha/2} \quad (14)$$

so a divergent local spin susceptibility requires a spin fluid with $d \leq 2$. Si et al are thus motivated to propose that the non-trivial physics of the heavy fermion QCP is driven by the formation of a two-dimensional spin fluid. Si et al consider an impurity spin within an effective medium in which the local Weiss field H has a critical power-spectrum (fig. 5.)

$$\langle |H(\omega)|^2 \rangle \equiv \chi_o^{-1}(\omega) = \omega^\gamma \quad (15)$$

where ϵ is self-consistently evaluated using a dynamical mean-field theory, where q -dependence of self-energies is dropped. In principle, the method solves the dynamical spin susceptibility of the impurity $\chi^{-1}(\omega) = \chi_o^{-1}(\omega) + M(\omega)$. This, in turn furnishes a “spin self-energy” $M(\omega)$ used to determine the spin susceptibility of the medium $\chi^{-1}(\vec{q}, \omega) = J(\vec{q}) + M(\omega)$.

Si et al find that a self-consistent solution is obtained for $\epsilon = 1$, if the spin-self energy contains a separate power-law dependence $M(\omega) \sim \omega^\alpha$ with an exponent $\alpha = 1/\rho\Gamma$ which is determined by the density of states

ρ and band-width Λ of the bond-strengths in the two-dimensional spin fluid. Although self-consistency requires a new power-law in the spin susceptibility, independent solutions of the impurity model have not yet shown that this feature is indeed generated by a critical Weiss field. This theory nevertheless raises many interesting questions:

1. Is the requirement of a two dimensional spin fluid consistent with experiment? For example- does the the cubic (and hence manifestly three dimensional) quantum critical material, $CeIn_3$ display a divergent specific heat?
2. If the spin-fluids are quasi-two dimensional, do we expect an ultimate cross-over to a three-dimensional QSDW scenario?
3. If α is non-universal, why are the critical exponents in $CeCu_{6-x}Au_x$ and $YbRh_2Si_{2-x}Ge_x$ so similar?
4. What stabilizes the local quantum criticality against intersite couplings?

2. Towards the critical Lagrangian.

If the heavy fermion QCP is a truly three-dimensional phenomenon, then a different approach is needed- we need to search for a new class of critical Lagrangian with $d_u > 3$ [41]. On general grounds, the existence of a Fermi liquid in the paramagnetic phase tells us that it must find expression in terms of the quasiparticle fields ψ in the Fermi liquid,

$$L = L_F[\psi] + L_{F-M}[\psi, M] + L_M[M]. \quad (16)$$

where L_F describes the heavy Fermi liquid, far from the magnetic instability, L_M describes the magnetic excitations that emerge above the energy scale $T_F^*(P)$.

L_{F-M} describes the way that the quasiparticles couple to and decay into critical magnetic modes; it also determines the type of transformation which takes place in the Fermi surface which takes at the QCP. This last point follows because away from the QCP, magnetic fluctuations can be ignored in the ground-state, so that $L_M \rightarrow 0$. In the paramagnetic phase, $\langle M \rangle = 0$ so $L_{FM} \rightarrow 0$, but in the antiferromagnetic phase $\langle M \rangle \neq 0$, i.e.

$$L_{\text{eff}} = \begin{cases} L_F^*[\psi] & \text{paramagnet} \\ L_F^*[\psi] + L_{FM}[\psi, \langle M \rangle] & \text{a.f.m.} \end{cases}$$

where the asterix denotes the finite renormalizations derived from zero-point fluctuations in the magnetization.

If the staggered magnetization is the fundamental critical field, then we are forced to couple the magnetic modes directly to the spin density of the Fermi liquid

$$L_{F-M}^{(1)} = g \sum_{\mathbf{k}, \mathbf{q}} \psi_{\mathbf{k}-\mathbf{q}}^\dagger \vec{\sigma} \psi_{\mathbf{k}} \cdot \vec{M}_{\mathbf{q}}. \quad (17)$$

But once the staggered magnetization condenses, this leads directly back to a static spin density wave (Fig 4).

An alternative possibility is suggested by the observation that the magnetism develops spinorial character in the heavy fermi liquid. The Luttinger sum rule [42] governing the Fermi surface volume V_{FS} “counts” both the electron density n_e and the number of the number of local moments per unit cell n_{spins} [43,44] :

$$2 \frac{V_{FS}}{(2\pi)^3} = n_e + n_{spins}. \quad (18)$$

The appearance of the spin density in the Luttinger sum rule reflects the composite nature of the heavy quasiparticles, formed from bound-states between local moments and high energy electron states. Suppose the spinorial character of the magnetic degrees of freedom seen in the paramagnet *also* manifests itself in the decay modes of the heavy quasiparticles. This would imply that at the QCP, the staggered magnetization factorizes into a spinorial degree of freedom $\vec{M}(x) = z^\dagger(x) \vec{\sigma} z(x)$, where z is a two-component spin 1/2 Bose field. “Spinorial magnetism” affords a direct coupling between the magnetic spinor z and the heavy electron quasi-particles via an inner product, over the spin indices

$$L_{F-M}^{(2)} = g \sum_{\mathbf{k}, \mathbf{q}} [z_{\mathbf{k}-\mathbf{q}\sigma}^\dagger \psi_{\mathbf{k}\sigma} \chi_{\mathbf{q}}^\dagger + \text{H.c.}], \quad (19)$$

where conservation of exchange statistics obliges us to introduce of a spinless charge e fermion χ . This would imply that the composite heavy electron decays into a neutral “spinon” and a spinless charge e fermion $e_\sigma^- \rightleftharpoons s_\sigma + \chi^-$. From this perspective, the heavy fermion QCP involves spin-charge separation, and the critical Lagrangian is a gauge theory.

To go beyond this general discussion we need to answer various questions:

- Can we connect the appearance of local criticality with an underlying gauge symmetry?
- What is the link between the heavy fermion QCP and metamagnetism?
- Is there a fundamental reason why the heavy fermion QCP exhibits H/T scaling (e.g as opposed to H^δ/T scaling) ?
- What feature in the critical Lagrangian can push the upper-critical dimension above 3?

On the experimental front, Hall constant measurements may provide a good way to discern between the spin density wave and composite quasiparticle alternative. In the former, regions around the hot-line do not

contribute to the Hall conductivity, and the change in the Hall constant is expected to evolve as the square of the staggered magnetization [8]. By contrast, the composite fermion scenario leads to a much more rapid evolution: depending on whether the density of spinless fermions is finite at the QCP the Hall constant which either jumps or evolves linearly with the staggered magnetization [8].

$$\Delta R_H \propto \begin{cases} M_{\mathbf{Q}}^2, & \text{(vectorial)} \\ O(1), M_{\mathbf{Q}} & \text{(spinorial)} \end{cases} \quad (20)$$

The only available Hall measurement at a QCP to date shows a change in sign takes place in the close vicinity of the QCP in critical $CeCu_{1-x}Au_x$, it is not yet clear whether there is a discontinuity at the transition [45]. This is clearly an area where more experimental input is highly desirable.

F. Summary

This paper has discussed the origin of the mass divergence at the heavy fermion quantum critical point, emphasizing that a quantum spin density wave picture can not explain the observed properties. The proposal of fundamentally new kinds of quantum critical point have been reviewed. This is clearly an area with a huge potential for progress both on the experimental, and theoretical front.

-
- [1] S. Sachdev, “*Quantum Phase Transitions*”, (Cambridge University Press, 1999).
 - [2] D. Obertelli, J. R. Cooper and J. L. Tallon, Phys. Rev. **B46**, 14928 (1992).
 - [3] C. Castellani, C. Di castro and M. Grilli, Z. Phys. B **103**, 137 (1997); J. Tallon et al, Phys. Stat. Solidi (b) 215, 531 (1999).
 - [4] L. D. Landau, Soviet Physics, JETP **3**, 920 (1957), (Zh. Eksp. Teor. Fiz. **30**, 1058 (1956)).
 - [5] D. Pines and P. Nozières, “The Theory of Quantum Liquids” (Benjamin, New York, 1966), vol 1.
 - [6] P. Nozières, “Theory of Interacting Fermi Systems”, (Benjamin, New York, 1964).
 - [7] C. M. Varma, Z. Nussinov and W. van Saarloos, “Singular Fermi Liquids”, submitted to Physics Reports, cond-mat/0103393.
 - [8] P. Coleman, C. Pépin, Qimiao Si and Revaz Ramazashvili, J. Phys. Cond. Matt **13** 273 (2001).
 - [9] O. Trovarelli, C. Geibel, S. Mederle, C. Langhammer, F.M. Grosche, P. Gegenwart, M. Lang, G. Sparn and F. Steglich, Phys. Rev. Lett. **85**, 626 (2000).
 - [10] H. v. Lohneysen et al. Phys. Rev Lett **72**, 3262 (1994).
 - [11] H. v. Lohneysen, J. Phys. Cond. Mat. **8**, 9689-9706 (1996).
 - [12] A. Schroeder et al., Nature **407**, 351(2000).
 - [13] K. Heuser et al., Phys. Rev. B **57**, R4198 (1998).
 - [14] M. Grosche et al., J. Phys. Cond Matt **12**, 533(2000).
 - [15] P. Gegenwart, F. Kromer, M. Lang, G. Sparn, C. Geibel and F. Steglich, Phys. Rev. Lett **82**, 1293 (1999).
 - [16] S.R. Julian et al., J. Phys. Cond. Mat. **8**, 9675 (1996).
 - [17] D. Braithwaite, T. Fukuhara, A. Demeur, I. Sheikin, S. Kambe, J-P. Brison, K. Maez, T. Naka and J. Floquet, J. Phys. Cond. Matt **12**, 1339 (2000).
 - [18] O. Trovarelli et al, to be published (2001).
 - [19] P. Estrela, A. de Visser, T. Naka, F.R. de Boer, L. C. J. Pereira, cond-mat/0106292, proceedings of SCES 2001.
 - [20] N. Mathur et al. Nature **394**, 39 (1998).
 - [21] P. Estrela, A. de Visser, F.R. de Boer, T. Naka and L. Shlyk, Phys. Rev. **B63**, 212409 (2001).
 - [22] P. Gegenwart et al. Phys. Rev. Lett. **81**, 1501 (1998).
 - [23] Y. Aoki et al, Journal of Magnetism and Magnetic Materials **177-181**, 271 (1998).
 - [24] J. S. Kim, D. Hall, K. Heuser and G. R. Stewart, Solid State Comm. **114**, 413 (2000).
 - [25] R. S. Perry et al., Phys. Rev. Lett. **86**, 2661 (2001).
 - [26] G. Knebel, D. Braithwaite, P. Canfield, G. Lapertot and J. Flouquet, to be published (2001).
 - [27] E/T scaling was first introduced as part of the Marginal Fermi Liquid phenomenology by C. M. Varma, P. B. Littlewood, S. Schmitt-Rink, E. Abrahams and A. E. Ruckenstein, Phys. Rev. Lett. **30**, 1996 (1989).
 - [28] E/T scaling was first observed in the disordered non-Fermi liquid material $UCu_{5-x}Pd_x$, M.C. Aronson, R. Osborn, R.A. Robinson, J.W. Lynn, R. Chau, C.L. Seaman, and M.B. Maple, Phys. Rev. Lett. **75**, 725 (1995).
 - [29] M. Blume, V. J. Emery and R. B. Griffiths, Phys. Rev. **4**, 1071, (1971).
 - [30] J. A. Hertz, Phys. Rev. B **14**, 1165 (1976).
 - [31] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, chap 30 p. 710 (Oxford University Press, 1993).
 - [32] S. Sachdev and J. Ye, Phys. Rev. Lett. **69**, 2411 (1992).
 - [33] M. T. Beal-Monod, Physica **109 & 110B**, 1837 (1982).
 - [34] T. Moriya and J. Kawabata, J. Phys. Soc. Japan **34**, 639 (1973); J. Phys. Soc. Japan **35**, 669 (1973).
 - [35] A. J. Millis, Phys. Rev. B **48**, 7183 (1993).
 - [36] O. Stockert, H.v. Lohneysen, A. Rosch, N. Pyka, M. Loewenhaupt, Phys. Rev. Lett. **80**, 5627 (1998).
 - [37] B. Fak, J. Flouquet, G. Lapertot, T. Fukuhara, H. Kadowaki, J. Phys. Cond. Matt **12**, 5423 (2000).
 - [38] A. Sengupta, Phys. Rev. B **62**, 4041, (2000).
 - [39] Q. Si, S. Rabello, K. Ingersent and J. L. Smith, cond-mat/0011477, to appear in Nature (2001).
 - [40] S. Doniach, Physica B **91**, 231 (1977).
 - [41] P. Coleman, C. Pépin, A. M. Tsvetlik, Phys. Rev. B, **62**, 3852-3868 (2000).
 - [42] J. M. Luttinger and J. C. Ward, Phys Rev. **118**, 1417 (1960), J. M. Luttinger, Phys. Rev. **119**, 1153 (1960).
 - [43] R. M. Martin, Phys. Rev. Lett. **48**, 362 (1982).
 - [44] M. Oshikawa, Phys. Rev. Lett. **84**, 3370 (2000).
 - [45] T. Nakimi, H. Sato, J. Jarakawa, H. Sugawara, Y. Aoki, R. Settai and Y. Onuki, Physica B **281-282**, 359 (2000).